

SIMPLE $SL(n)$ -MODULES WITH NORMAL CLOSURES OF MAXIMAL TORUS ORBITS

K. KUYUMZHIYAN

ABSTRACT. Let T be the subgroup of diagonal matrices in the group $SL(n)$. The aim of this paper is to find all finite-dimensional simple rational $SL(n)$ -modules V with the following property: for each point $v \in V$ the closure $\overline{Tv}$ of its T -orbit is a normal affine variety. Moreover, for any $SL(n)$ -module without this property a T -orbit with non-normal closure is constructed. The proof is purely combinatorial: it deals with the set of weights of simple $SL(n)$ -modules. The saturation property is checked for each subset in the set of weights.

INTRODUCTION

Let T be an algebraic torus defined over an algebraically closed field $\mathbb{k}$ of characteristic zero. Recall that an irreducible algebraic T -variety X is called *toric* if X is normal and T acts on X with an open orbit. This class of varieties plays an important role in algebraic geometry, topology and combinatorics due to its remarkable description in terms of convex geometry, see [Ful]. Assume that the torus T acts on a variety Y . Then the closure $X = \overline{T y}$ of the T -orbit of a point $y \in Y$ is a natural candidate to be a toric variety. To verify it, one should check that X is normal.

During last decades, normality of torus orbits' closures was an object of numerous investigations. For example, let G be a semisimple algebraic group with a Borel subgroup B and a maximal torus $T \subset B$. In [Kl], it was proved that the closure of a general T -orbit on the flag variety G/B is normal. Later it was shown that the closure of a general T -orbit in G/P , where $P \subset G$ is a parabolic subgroup, is also normal, see [Da]. Examples of non-normal closures of non-general torus orbits can be found in [CK].

Now let us consider a finite-dimensional rational T -module V . There exists an easy combinatorial criterion of normality of $\overline{T v}$ for a vector $v \in V$. Namely, let $v = v_{\chi_1} + \dots + v_{\chi_m}$, $v_{\chi_i} \neq 0$, be the weight decomposition of the vector v . Consider the corresponding set of T -weights $\chi_1, \dots, \chi_m$. If we take $\chi_1, \dots, \chi_m$ as elements of the character lattice $\mathfrak{X}(T)$, we can generate a semigroup $\mathbb{Z}_+(\chi_1, \dots, \chi_m)$, a sublattice $\mathbb{Z}(\chi_1, \dots, \chi_m)$, and a rational polyhedral cone $\mathbb{Q}_+(\chi_1, \dots, \chi_m)$. The set $\chi_1, \dots, \chi_m$ is called *saturated* if $\mathbb{Z}_+(\chi_1, \dots, \chi_m) = \mathbb{Z}(\chi_1, \dots, \chi_m) \cap \mathbb{Q}_+(\chi_1, \dots, \chi_m)$. It is well known (see [KKMSD, page 5]) that the following two conditions are equivalent: the set $\{\chi_1, \dots, \chi_m\}$ is saturated and the closure $\overline{T v}$ of the T -orbit $T v$ is normal. There is an analogous criterion for the T -action on the projectivisation $\mathbb{P}(V)$, see [CK].

The saturation property occurs in many algebraic and geometric problems. In [Wh], it was proved that the set of incidence vectors of the bases of a realizable matroid is saturated. The geometric conclusion of this fact is that for any point y in the affine cone over the classical Grassmannian $\text{Gr}(k, n)$ the closure $\overline{T y}$ is normal.

Taken a finite graph Γ with n vertices, one can associate a finite collection $M(\Gamma)$ of vectors in the lattice $\mathbb{Z}^n$ with it:

$$M(\Gamma) = \{\varepsilon_i + \varepsilon_j \mid (ij) \text{ is an edge of } \Gamma\},$$

where $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ is the standard basis of $\mathbb{Z}^n$. The saturation property for this set is equivalent to the fact that for arbitrary two minimal odd cycles C and C' in Γ , either C and C' have a common vertex or there exists an edge of Γ joining a vertex of C with a vertex of C' (see [OH] and [SVV]). Algebraically, the saturation property for $M(\Gamma)$ is equivalent to the integral closeness for the subalgebra $\mathcal{A}(\Gamma)$ of the polynomial algebra $\mathbb{k}[x_1, x_2, \dots, x_n]$,

$$\mathcal{A}(\Gamma) = \mathbb{k}[x_i x_j \mid (ij) \text{ is an edge of } \Gamma]$$

in its field of fractions $Q\mathcal{A}(\Gamma)$.

Some general results concerning quivers and the saturation property were obtained in [Chi]. It was shown that a finite, connected quiver Q without oriented cycles is a Dynkin or Euclidean quiver if and only if all orbit semigroups of representations of Q are saturated.

In the paper [Mo], the following problem is solved. Let G be a semisimple algebraic group with a maximal torus T and V be its adjoint module. For which G for all $v \in V$ the closure $\overline{Tv}$ is normal? The surprising fact is that for $G = SL(n)$ this is always the case (see also [Stu1, Ex. 3.7], [Stu2], [Mo] and [BZ, Prop.2.1]). In [BZ], this combinatorial result is interpreted in terms of representations of quivers.

The aim of this paper is to classify all simple finite-dimensional rational $SL(n)$ -modules V such that for any $v \in V$ the closure $\overline{Tv}$ is normal.

Main Theorem. *The representations below, together with their dual, form the list of all irreducible representations of $SL(n)$ where all maximal torus orbits' closures are normal:*

- (1) *the tautological representation of $SL(n)$;*
- (2) *the adjoint representation of $SL(n)$;*
- (3) *exceptional cases:*

<i>Group</i>	<i>Highest weight</i>	<i>G-module</i>
$SL(2)$	$3\pi_1$	$S^3\mathbb{k}^2$
$SL(2)$	$4\pi_1$	$S^4\mathbb{k}^2$
$SL(3)$	$2\pi_1$	$S^2\mathbb{k}^3$
$SL(4)$	π_2	$\Lambda^2\mathbb{k}^4$
$SL(5)$	π_2	$\Lambda^2\mathbb{k}^5$
$SL(6)$	π_2	$\Lambda^2\mathbb{k}^6$
$SL(6)$	π_3	$\Lambda^3\mathbb{k}^6$

The paper is organized as follows. In Section 1 we give some algebraic definitions and reformulate the problem in combinatorial terms. From that point, it remains to check the saturation property for any subset in the system of T -weights of a simple $SL(n)$ -module. In Section 2 we prove that the saturation property holds for each subset in the set of weights of the representations listed in the Main Theorem. We also give a new proof of theorem [Mo, Thm.1]. It is the most non-trivial positive case, where the dimension of V is not bounded. When possible, reasoning uses the graph theory language. Our reference for graph theory

is [Ha]. In Section 3 we produce non-saturated subsets in sets of weights for all other representations. If the set of weights of the representation with the highest weight λ is a subset in the set of weights of the representation with the highest weight μ , and a non-saturated subset for λ is known, then one can use it as a non-saturated subset for μ . Fundamental representations form the most difficult case. To work with them, we use the following observation. If a non-saturated subset in the set of weights of the k th fundamental representation of $SL(n)$ is found, then the analogous non-saturated subset exists in the set of weights of the k th fundamental representation of $SL(n+k)$.

Some auxiliary results used in the proof may be of independent interest.

Lemma (Reformulation of Lemma 2.5). *Let $M_6 = \{(\varepsilon_1, \dots, \varepsilon_6) \mid \varepsilon_i = \pm 1, \sum \varepsilon_i = 0\}$ be the set of points in $\mathbb{R}^6$. Then each subset of this set is saturated.*

In further publications, we plan to give a classification of simple G -modules with normal T -orbit closures for other simple algebraic groups G .

The author is grateful to her scientific supervisor I.V. Arzhantsev for the formulation of the problem and fruitful discussions. Thanks are also due to I.I. Bogdanov for useful comments.

1. ALGEBRAIC BACKGROUND AND NOTATION

Let V be a finite-dimensional rational T -module. Given any character χ from the character lattice $\mathfrak{X}(T)$, define a weight subspace V_χ as $V_\chi = \{v \in V \mid t \cdot v = \chi(t)v\}$. It is well known that $V = \bigoplus_{\chi \in \mathfrak{X}(T)} V_\chi$, and only finitely many V_χ are nonzero. The set $\{\chi_1, \chi_2, \dots, \chi_k\}$ of those χ_i for which $V_{\chi_i} \neq 0$ is called *the system of weights* of the T -module V .

Let $\mathbb{Z}_+$ and $\mathbb{Q}_+$ denote the sets of integer and rational non-negative numbers, respectively; and let $v_1, v_2, \dots, v_m \in \mathbb{Q}^n$. Consider the semigroup $\mathbb{Z}_+(v_1, v_2, \dots, v_m) = \{n_1 v_1 + n_2 v_2 + \dots + n_m v_m \mid n_i \in \mathbb{Z}_+\} \subseteq \mathfrak{X}(T)$, the sublattice $\mathbb{Z}(v_1, v_2, \dots, v_m) = \{z_1 v_1 + z_2 v_2 + \dots + z_m v_m \mid z_i \in \mathbb{Z}\} \subseteq \mathfrak{X}(T)$, and the rational polyhedral cone $\mathbb{Q}_+(v_1, v_2, \dots, v_m) = \{q_1 v_1 + q_2 v_2 + \dots + q_m v_m \mid q_i \in \mathbb{Q}_+\} \subseteq \mathfrak{X}(T) \otimes_{\mathbb{Z}} \mathbb{Q}$. Define the following important property of the set $\{v_1, v_2, \dots, v_m\}$.

Definition. The set of points $\{v_1, v_2, \dots, v_m\} \subset \mathbb{Q}^n$ is called *saturated* if

$$\mathbb{Z}_+(v_1, v_2, \dots, v_m) = \mathbb{Z}(v_1, v_2, \dots, v_m) \cap \mathbb{Q}_+(v_1, v_2, \dots, v_m).$$

The following result provides a well-known combinatorial criterion of normality of the torus orbit closure, see [KKMSD]:

Theorem 1.1. *Consider a rational linear action of a torus T on a vector space V . Let $v = v_{\lambda_1} + \dots + v_{\lambda_s}$, $v_{\lambda_i} \neq 0$, be its weight decomposition. Then the closure $\overline{Tv}$ is normal if and only if the set of characters $\{\lambda_1, \dots, \lambda_s\}$ is saturated.*

Corollary 1.2. *Given a rational linear action of torus T on a vector space V ; let $\{\lambda_1, \dots, \lambda_s\}$ be the set of weights of this action. Then the closure $\overline{Tv}$ is normal for each $v \in V$ if and only if each subset in $\{\lambda_1, \dots, \lambda_s\}$ is saturated.*

Remark. The weight system is multiplied by -1 while changing a representation V of torus T with its dual representation. Hence the property of normality of all T -orbits is preserved.

Let $G = SL(n)$. We fix a maximal torus $T \subset G$ consisting of all diagonal matrices. An element $a = (a_1, a_2, \dots, a_n)$ of the lattice $\mathbb{Z}^n$ can be interpreted as a character χ_a of the torus T

in the following way: $\chi_a(t) = t_1^{a_1} t_2^{a_2} \dots t_n^{a_n}$, where $t = \text{diag}(t_1, t_2, \dots, t_n)$. Since $t_1 t_2 \dots t_n = 1$, the points a and b define the same character if and only if $a - b = \alpha(1, 1, \dots, 1)$. Each a from $\mathbb{Z}^n$ has a unique representation $a = \tilde{a} + \alpha(1, 1, \dots, 1)$, where $\tilde{a} \in \mathbb{Q}^n$, $\alpha \in \mathbb{Q}$, and $\sum \tilde{a}_i = 0$.

Let $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ be the standard basis of the lattice $\mathbb{Z}^n$, and

$$e_i = \tilde{\varepsilon}_i = \left(-\frac{1}{n}, -\frac{1}{n}, \dots, -\frac{1}{n}, \underset{i\text{th place}}{\frac{n-1}{n}}, -\frac{1}{n}, \dots, -\frac{1}{n} \right).$$

Notice that $e_1, e_2, \dots, e_n$ (further referred to as a *quasi-basis*) satisfy the only linear relation

$$e_1 + e_2 + \dots + e_n = 0. \quad (*)$$

Identify $\mathfrak{X}(T)$ with the $\mathbb{Z}$ -lattice generated by $e_1, e_2, \dots, e_n$. Recall that a weight χ_a is called *dominant* if and only if $a_1 \geq a_2 \geq \dots \geq a_n$. The *root lattice* for $SL(n)$ is a lattice generated by the vectors $e_1 - e_2, e_2 - e_3, \dots, e_{n-1} - e_n$. Due to the ambiguity of notation,

$$\Phi = \{a_1 e_1 + a_2 e_2 + \dots + a_n e_n \mid a_1 + a_2 + \dots + a_n \in \mathbb{Z}\}.$$

For a positive integer $s \mid n$ (i.e. $n = ss', s' \in \mathbb{Z}$), define

$$\mathbb{Z}_{\equiv 0(s)}(e_1, \dots, e_n) = \left\{ \sum_{i=1}^n x_i e_i \mid x_i \in \mathbb{Z}, \sum_{i=1}^n x_i \equiv 0 \pmod{s} \right\}.$$

In this notation the root lattice Φ coincides with $\mathbb{Z}_{\equiv 0(n)}(e_1, \dots, e_n)$.

Let V be a finite-dimensional simple rational $SL(n)$ -module, $M(V)$ be the system of weights of the module V with respect to the restricted action $T: V$. Introduce a partial order on $M(V)$: $\mu \succeq \nu$ if and only if for $\xi = \mu - \nu$ the following conditions hold: $\xi_1 \geq 0$, $\xi_1 + \xi_2 \geq 0$, $\dots$, $\xi_1 + \xi_2 + \dots + \xi_{n-1} \geq 0$. It is well known that $M(V)$ contains the only maximal element λ with respect to $\succeq$, it is called *the highest weight* of the module. The weight λ is dominant, moreover, for any dominant weight $\lambda \in \mathfrak{X}(T)$ there exists a unique simple $SL(n)$ -module $V(\lambda)$ with the highest weight λ (see [Hu, §20, §21]). The role of the Weyl group W is played here by the permutation group S_n , which acts on $\mathbb{Z}^n$ by permutations of coordinates. It is well known (see [Hu, §13, §21] or [VO, Chapter 4]) that

$$M(\lambda) := M(V(\lambda)) = \text{conv}\{w\lambda \mid w \in W\} \cap (\lambda + \Phi),$$

where $\text{conv}(M)$ denotes the convex hull of the set $M \subset \mathbb{R}^n$.

In our situation, Corollary 1.2 can be reformulated in the following way:

Proposition 1.3. *Let $V(\lambda)$ be a simple module of a semisimple group G with the highest weight λ . Then the closure of each T -orbit in $V(\lambda)$ is normal if and only if each subset in $M(\lambda)$ is saturated.*

2. POSITIVE RESULTS

In this section we prove that certain sets of weights are saturated. We use the following lemmas, their proof can be found in [Mo].

Lemma 2.1. *Let M be a non-saturated set and α be a vector such that $\alpha \in M$ and $-\alpha \in M$. Then either $M \setminus \{\alpha\}$ or $M \setminus \{-\alpha\}$ is non-saturated.*

Lemma 2.2. *Any set of linearly independent vectors is saturated.*

Lemma 2.3. *Let $v = q_1v_1 + \dots + q_mv_m$, where v, v_i are arbitrary vectors, and $q_i \in \mathbb{Q}_+$. Then one can choose a linearly independent subset $\{v_{i_1}, \dots, v_{i_s}\} \subset \{v_1, \dots, v_m\}$ and numbers $q'_{i_1}, \dots, q'_{i_s} \in \mathbb{Q}_+$ such that*

$$v = q'_{i_1}v_{i_1} + \dots + q'_{i_s}v_{i_s}.$$

2.1. The tautological representation. The highest weight of the tautological representation equals e_1 . One should prove that each subset in $\{e_1, \dots, e_n\}$ is saturated. Using Lemma 2.2, we obtain that any proper subset of this set is saturated. It is easy to see that this set itself is also saturated because if one takes a $\mathbb{Z}$ -combination of vectors e_i , then it becomes a $\mathbb{Z}_+$ -combination after adding $(*)$ with a positive coefficient.

2.2. The adjoint representation. Its highest weight λ is equal to $e_1 - e_n$. Acting by $W = S_n$, we get all vectors of the form $e_i - e_j$. Taking the convex hull adds only $\bar{0}$ to this set. We yield $M(\lambda) = \{0\} \cup \{e_i - e_j \mid 1 \leq i, j \leq n\}$. The saturation property for this set was proved, e.g., in [Mo, Thm.1]; nevertheless, we present another proof based on a graph theory approach.

Theorem 2.4. *Any subset in $M(\lambda) = \{0\} \cup \{e_i - e_j \mid 1 \leq i, j \leq n\}$ is saturated.*

Proof. Fix a nonempty subset $S \subseteq M(\lambda)$. Construct a directed graph Γ according to S . Let Γ have n vertices $A_1, A_2, \dots, A_n$, where A_i corresponds to e_i . An arc A_iA_j exists for each element $e_i - e_j$ from S . Taken a formal linear combination $q_1v_1 + q_2v_2 + \dots + q_nv_n$, where $q_i \in \mathbb{Q}$ and v_i are the arcs of Γ , one can get a corresponding linear combination $p_1e_1 + p_2e_2 + \dots + p_ne_n$ by substituting instead of all v_i the corresponding vectors $e_j - e_k$. If the coefficients q_i are given, one can write the explicit formula for p_i . Namely,

$$p_i = \sum_{\substack{v_l = A_iA_m \\ \text{is an arc of } \Gamma}} q_l - \sum_{\substack{v_l = A_mA_i \\ \text{is an arc of } \Gamma}} q_l. \quad (1)$$

Our aim is to prove that S is saturated. Reformulate the saturation property in terms of graphs. We have to show that if $v_1, v_2, \dots, v_r$ is the set of edges of Γ , then each vector v from $\mathbb{Q}_+(v_1, v_2, \dots, v_r) \cap \mathbb{Z}(v_1, v_2, \dots, v_r)$ is also an element of $\mathbb{Z}_+(v_1, v_2, \dots, v_r)$. Take a vector $v \in \mathbb{Q}_+(v_1, v_2, \dots, v_r) \cap \mathbb{Z}(v_1, v_2, \dots, v_r)$. We can multiply all its coordinates by an integer m in such a way that all coefficients of the corresponding $\mathbb{Q}_+$ -combination become integer, i.e. $v_1 = mv \in \mathbb{Z}_+(v_1, v_2, \dots, v_r) \cap m\mathbb{Z}(v_1, v_2, \dots, v_r)$, where $m\mathbb{Z}(v_1, v_2, \dots, v_r)$ denote the set of linear combinations of v_i where all coefficients are divisible by m . Since $v_1 \in m\mathbb{Z}(v_1, v_2, \dots, v_r)$, we can apply (1) for the corresponding $m\mathbb{Z}$ -combination and obtain that all its coordinates are divisible by m . Now consider the corresponding $\mathbb{Z}_+$ -combination for v_1 . If all vectors from Γ enter with multiplicities $:m$, then we are done: we can divide them by m and get the required combination for v . Otherwise, take all v_i entering into the $\mathbb{Z}_+$ -combination for v_1 with multiplicities $\not:m$ and name them *bad*. Let Γ' be the subgraph in Γ formed by all bad edges. If we count the sum

$$p_i = \sum_{\substack{v_l = A_iA_m \\ \text{is an arc of } \Gamma'}} q_l - \sum_{\substack{v_l = A_mA_i \\ \text{is an arc of } \Gamma'}} q_l \quad (2)$$

at a vertex A_i , then it will be divisible by m since we excluded only the edges with multiplicities $\not:m$. This means that Γ' cannot have terminal vertices. Now find a cycle in Γ' (not

necessarily oriented!). At each step, we will increase by 1 the values at the edges of this cycle oriented clockwise and decrease by 1 at the edges oriented counter-clockwise. This operation does not change v and v_1 . In several steps (not more than m) the value at an edge of Γ' will be divisible by m . So the number of bad edges has decreased. Repeating this operation, we will change the values at the edges in such a way that they all will become divisible by m , and the vector v_1 (and, also, v) will not change. Dividing all the coefficients of the constructed $\mathbb{Z}_+$ -combination by m , we get the required $\mathbb{Z}_+$ -combination for v . $\square$

Definition. We will mean by *NSS* a non-saturated subset $\{v_1, v_2, \dots, v_s\}$ in the set M of weights of a representation. By *ENSS* we will mean the NSS together with a vector v , where $v \in \mathbb{Z}(v_1, v_2, \dots, v_m) \cap \mathbb{Q}_+(v_1, v_2, \dots, v_m)$, and $v \notin \mathbb{Z}_+(v_1, v_2, \dots, v_m)$.

2.3. The representation of $SL(2)$ with the highest weight $3\pi_1$. One has to verify the saturation property for each subset in the set $M = \{(\frac{3}{2}, -\frac{3}{2}), (\frac{1}{2}, -\frac{1}{2}), (-\frac{1}{2}, \frac{1}{2}), (-\frac{3}{2}, \frac{3}{2})\}$ (in the usual basis). If there exists an NSS, then Lemma 2.1 is applicable, and this NSS can be reduced either to $\{(\frac{3}{2}, -\frac{3}{2}), (\frac{1}{2}, -\frac{1}{2})\}$ or to $\{(\frac{3}{2}, -\frac{3}{2}), (-\frac{1}{2}, \frac{1}{2})\}$ (up to sign change). But these subsets are both saturated. This means that the initial NSS is also saturated, a contradiction.

2.4. The representation of $SL(2)$ with the highest weight $4\pi_1$. Apply Lemma 2.1 to the set $M(\lambda) = \{(2, -2), (1, -1), (0, 0), (-1, 1), (-2, 2)\}$. If an NSS exists, then it contains not more than one vector of $(1, -1), (-1, 1)$, and not more than one of $(2, -2), (-2, 2)$. So, up to sign change, the NSS coincides with the set $\{(2, -2), (1, -1)\}$ or with the set $\{(2, -2), (-1, 1)\}$. But they are both saturated. We get a contradiction.

2.5. The representations of $SL(4)$, $SL(5)$ and $SL(6)$ with the highest weight π_2 . The set of weights $M(\lambda)$ is equal to $\{e_i + e_j \mid 1 \leq i, j \leq n, i \neq j\}$, where $n = 4, 5, 6$. A graph Γ with n vertices can be associated with any subset $S \subseteq M(\lambda)$: an edge connecting i th and j th vertices exists whenever $e_i + e_j \in S$. Suppose that there exists an ENSS $\{w; v_1, \dots, v_m \mid v_i \in M(\lambda)\}$:

$$w = z_1 v_1 + \dots + z_m v_m = q_1 v_1 + \dots + q_m v_m, \quad z_i \in \mathbb{Z}, q_i \in \mathbb{Q}_+, \\ w \notin \mathbb{Z}_+(v_1, \dots, v_m).$$

Consider all v_i occurring into the right hand of this equality with a nonzero coefficient q_i . By Lemma 2.3, we may assume that they are linearly independent. To simplify the reasoning, consider vector $v = w - [q_1]v_1 - \dots - [q_m]v_m$ instead of w . It is easy to see that v belongs to $\mathbb{Z}(v_1, \dots, v_m)$, to $\mathbb{Q}_+(v_1, \dots, v_m)$ and does not belong to $\mathbb{Z}_+(v_1, \dots, v_m)$. We yield that $\{v; v_1, \dots, v_m\}$ is also an ENSS. After this change all the coefficients of the $\mathbb{Q}_+$ -combination belong to $[0, 1)$.

Construct a subgraph $\Gamma' \subset \Gamma$: take all the vertices of Γ and all the edges of Γ entering into the $\mathbb{Q}_+$ -combination above with nonzero coefficients. Write the coefficients of the $\mathbb{Q}_+$ -combination at the edges of Γ' . The further proof consists of a search of all possible graphs Γ' . The following observations will simplify the search.

(0.1) The number of edges in each connected component of Γ' is not greater than the number of vertices (if not, the vectors corresponding to the edges of this component will be linearly dependent).

(0.2) The number of edges in Γ' is less than the number of vertices (it follows from $(*)$ that the dimension of the enveloping space equals $n - 1$).

(0.3) Graph Γ' does not contain even cycles. It follows from the fact that the edges of an even cycle are linearly dependent: their alternating sum is null.

(0.4) It follows from (0.1) and (0.3) that each connected component of Γ' either is a tree or contains exactly one cycle. In the second case this cycle is always odd.

(0.5) It follows from (0.2) that Γ' has a vertex of degree 0 or 1.

At each vertex, count the sum of all coefficients on the incident edges, then for each sum take its fractional part. All these fractional parts are equal due to the fact that all the sums in vertices (they equal the coordinates of v) become integer after subtracting $(*)$ with a proper coefficient. Now we conclude that

(0.6) Γ' does not contain vertices of degree 0 and 1 simultaneously: if it does, the fractional parts of the sums in vertices are all equal to 0, but in the terminal vertex this sum has only one summand and is not an integer.

We consider these two cases independently.

Case 1. Graph Γ' has a vertex of degree 0.

(1.1) Any other connected component of this graph is either a point or has no terminal vertices (it follows from (0.6)). Moreover, it follows from (0.4) that it is an odd cycle.

(1.2) We have $n \leq 6$, consequently, the number of edges in Γ' is ≤ 5 , but any odd cycle has ≥ 3 edges, and we yield that Γ' has at most 1 cycle.


Fulfill an exhaustive search within all graphs Γ' having a vertex of degree 0.

$$\begin{aligned} n = 4, & \quad \text{graph is a cycle of length 3,} \\ n = 5, & \quad \text{graph is a cycle of length 3,} \\ n = 6, & \quad \text{graph is a cycle of length 3,} \\ n = 6, & \quad \text{graph is a cycle of length 5.} \end{aligned}$$

The only possible $\mathbb{Q}_+$ -combination in these cases is $\frac{1}{2}(v_1 + \dots + v_s)$. This means that v is a sum of e_i corresponding to the vertices of the cycle. But it does not lie in $\mathbb{Z}(v_1, \dots, v_m)$ when n is even. When $n = 5$, consider also the graph Γ . Since v is a $\mathbb{Z}$ -combination of the edges of Γ , Γ has more than 3 edges: $\Gamma \supset \Gamma'$, $\Gamma \neq \Gamma'$ and Γ' has 3 edges. In the representation above the sum of coefficients of v is odd, hence we must apply $(*)$ to the existing $\mathbb{Z}$ -combination to get the same representation. For this purpose the edges from $\Gamma \setminus \Gamma'$ should touch all the vertices of Γ (we name this property $(**)$).

In fig. 1 the graph Γ' is drawn. To satisfy $(**)$, Γ must contain at least the following edges (up to symmetry): see fig. 2, 3 or 4. The vertices corresponding to e_i are called V_i . But in all cases we get a contradiction since $e_1 + e_2 + e_3$ is already a $\mathbb{Z}_+$ -combination:

$$\begin{aligned} \text{in Fig. 2} \quad & e_1 + e_2 + e_3 = V_1V_2 + V_2V_3 + V_1V_3 + V_4V_5, \\ \text{in Fig. 3} \quad & e_1 + e_2 + e_3 = V_4V_2 + V_2V_1 + V_1V_3 + V_3V_5, \\ \text{in Fig. 4} \quad & e_1 + e_2 + e_3 = V_4V_2 + V_2V_5 + 2V_1V_3. \end{aligned}$$



We have shown that the counterexample does not exist in the case when Γ' has a vertex of degree 0.

Case 2. Graph Γ' has a vertex X of degree 1. Let XY be an edge incident to X . We need to subtract $(*)$ with the same multiplicity as is at XY . We yield

(2.1) since X is a terminal vertex of Γ' , either XY is a connected component of Γ' or the vertex degree of Y is ≥ 3 . Indeed, suppose that the vertex degree of Y is 2. Let YZ be the second edge incident to Y , and let q be the value written at YZ . Then after subtracting $(*)$ the coefficient at Y becomes equal to q , but it must be integer, and we know that $q \in (0, 1)$. This is a contradiction.

Find all possible connected components of Γ' .

On 2 vertices:

On 3 vertices:

On 4 vertices: and

Notice that if we have a connected component of Γ' on 5 or 6 vertices, it is the only connected component of Γ' . Using this and (0.2), we obtain that Γ' is a tree. Apply (2.1). We have to consider only the following trees:

On 5 vertices:

On 6 vertices: and

But the edges of are linearly dependent (when $n = 6$, one should sum all the thin edges, then subtract the thick one, and obtain $(*)$). Therefore, this graph should not be considered.

Fulfill an exhaustive search within all graphs Γ' on n vertices satisfying all the conditions above. In the case when one of the connected components of Γ' is a claw (i.e., all the edges are incident to one vertex), its central vertex will correspond to e_1 (it is easy to see that Γ' cannot have more than one claw).

n	Splitting into connected components	Permissible graphs
4	$2 + 2$	
4	4	or
5	$2 + 3$	
5	5	
6	$2 + 2 + 2$	
6	$2 + 4$	or
6	6	

The graphs $\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}$ and $\begin{smallmatrix} \bullet \\ \diagup \end{smallmatrix}$ do not satisfy our conditions: their edges are linearly dependent.

If we start with $\begin{smallmatrix} \bullet \\ \diagup \end{smallmatrix}$, we can obtain only e_1 as the $\mathbb{Q}_+$ -combination: all the three edges must appear in the $\mathbb{Q}_+$ -combination with the same coefficient, let a , $a \in (0, 1)$. We sum these three vectors, obtain $3ae_1 + ae_2 + ae_3 + ae_4$, and subtract $(*)$ with a necessary coefficient. Finally we obtain $2ae_1$. In this notation it already has integer coordinates (equal to zero), this means that all the other coordinates, $2a$ among them, must be integers, $a = \frac{1}{2}$, $v = e_1$. But v cannot be obtained as the $\mathbb{Z}$ -combination of the vectors of the type $e_i + e_j$. Indeed, each v_i has an even sum of coordinates, n is even, subtracting $(*)$ with an integer coefficient does not change parity of the sum of coordinates, this proves that any vector from $\mathbb{Z}_+\{v_i\}_{i=1}^m$ has an even sum of coordinates.

The edges of graph $\begin{smallmatrix} \bullet \\ \diagup \end{smallmatrix}$ are linearly dependent (here $n = 5$) because $(2 \cdot \text{first edge} + \text{the sum of the edges of the cycle}) = 0$.

Graph $\begin{smallmatrix} \bullet \\ \diagup \end{smallmatrix}$: using similar reasoning, $v = e_1$ or $2e_1$. But there exists an edge $\in \Gamma \setminus \Gamma'$, hence e_1 is a $\mathbb{Z}_+$ -combination of the edges of Γ : take the sum of thick edges of $\begin{smallmatrix} \bullet \\ \diagup \end{smallmatrix}$.


The edges of $\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}$ and $\begin{smallmatrix} \bullet \\ \diagup \end{smallmatrix}$ are linearly dependent.

In graph $\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}$ v may be equal only to e_1 , but e_1 can not be obtained as a $\mathbb{Z}$ -combination: 6 is even, the sum of coordinates of e_1 is odd.

In graph $\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}$ vector v has to be proportional to e_1 , moreover, the coefficient must be even (we use the reasoning as above, from the fact that 6 is even it follows that the sum of coordinates is even for any vector from $\mathbb{Z}(v_1, \dots, v_m)$). But if we add any edge to this set, $2e_1$ will be obtained as a $\mathbb{Z}_+$ -combination: take the sum of thick edges of $\begin{smallmatrix} \bullet \\ \bullet \end{smallmatrix}$.

All the cases are considered, this completes the proof.

2.6. The representation of $SL(3)$ with the highest weight $2\pi_1$. Its highest weight λ is equal to $2\pi_1 = 2e_1$, and all the weights of this representation are pointed in the figure below.



Assume that this set contains an NSS $\{v_1, \dots, v_m\}$ (and ENSS $\{v; v_1, \dots, v_m\}$). Consider the following possibilities.

If this NSS contains both $-e_1$ and $-e_2$, then it does not contain $-e_3$. Indeed, if it does, then $\mathbb{Z}_+(-e_1, -e_2, -e_3) = \mathbb{Z}(-e_1, -e_2, -e_3)$, and this NSS cannot be non-saturated. Using the similar reasoning, we get that it contains at most one of the vectors $2e_1$ and $2e_2$: otherwise $\mathbb{Z}_+(-e_1, -e_2, 2e_1, 2e_2) = \mathbb{Z}(-e_1, -e_2, -e_3)$, and this NSS also cannot be non-saturated. Consequently, this NSS coincides (up to the indices renumbering) either with $(-e_1, -e_2, 2e_1)$ or with $(-e_1, -e_2, 2e_1, 2e_3)$. But these subsets are saturated.

If the NSS has no vectors of form $-e_i$, then $\{v_1, \dots, v_m\} \subset \{2e_1, 2e_2, 2e_3\}$. But in this case the NSS is also saturated.

If the NSS has exactly one vector of form $-e_i$, suppose $-e_1$, then fix a representation $n_1e_1 + n_2e_2 + n_3e_3$ for v , $n_1, n_2, n_3 \in \mathbb{Z}$. Then n_2 and n_3 have the same parity, since e_2 and e_3 occur in the corresponding $\mathbb{Z}$ -combination for v only as $2e_2$ and $2e_3$. Apply Lemma 2.3.

We get a representation $v = q_1v_1 + q_2v_2$, where v_1 and v_2 are linearly independent. We may suppose $q_i \in [0, 1)$ because otherwise v can be changed to a vector $v - v_i$, which will diminish (in some sense) the NSS. So the required combination will be either $\frac{2e_1}{2}$ or $\frac{2e_2+2e_3}{2}$. In the first case, recall that the NSS contains $-e_1$, and we yield e_1 as the $\mathbb{Z}_+$ -combination $2e_1 - e_1$. In the second case $-e_1$ is already the $\mathbb{Z}$ -combination, this is also a contradiction.

2.7. The representation of $SL(6)$ with the highest weight π_3 . The highest weight λ equals $e_1 + e_2 + e_3$, $M(\lambda) = \{e_i + e_j + e_k \mid 1 \leq i < j < k \leq 6\}$.

Lemma 2.5. *In the notation above, for any 5 linearly independent vectors $v_1, \dots, v_5 \in M(\lambda)$ the following equality holds:*

$$\mathbb{Z}(v_1, \dots, v_5) = \mathbb{Z}_{\equiv 0(3)}(e_1, \dots, e_6).$$

This means the following. For any vector $\mathbb{Z}(v_1, \dots, v_5)$, its sum of coordinates is $\equiv 3$, and this property does not depend on its representation. A surprising fact is that the reverse statement is true – if we take an integer vector with the sum of coordinates divisible by 3, then the vector with the same coordinates in quasi-basis can be obtained as a $\mathbb{Z}$ -combination of vectors $v_1, \dots, v_5$.

Proof. First, we are going to show that vector $(1, 1, 1, 0, 0, 0)$ lies in $\mathbb{Z}(v_1, \dots, v_5)$. Notice that if we take $-v_i$ instead of v_i for an index i , then the $\mathbb{Z}$ -lattice will not change. In quasi-basis it means that we take a vector with the complementary set of indices. Consequently, we may suppose that each v_i contains e_1 . Construct a graph with 5 vertices and 5 edges: each vertex corresponds to one of the integers 2, 3, 4, 5, 6, vertices i and j are connected with an edge if and only if there exists k , such that $v_k = e_1 + e_i + e_j$. Examine all possible graphs. Vectors $\{v_i\}_{i=1}^5$ are linearly independent by the data, this fact has the following conclusions:

- (1) the graph has no cycles of length 4 – otherwise we have a zero sum of form $v_1 - v_2 + v_3 - v_4$;
- (2) the graph has no vertices of degree 0 – otherwise a subgraph containing other 4 vertices has 5 edges, this means that it has a cycle of length 4.
- (3) this graph is connected. Indeed, if it has 2 or more connected components, none of which is a single vertex, then it has two connected components of 2 and 3 vertices respectively, which gives at most $3 + 1 = 4$ edges. This graph has a cycle since it has a sufficient number of edges. Using (1), we get that this cycle has 3 or 5 vertices, which means that it is odd.

For any vertex X of this graph there exists an odd cycle (maybe not a circuit) passing through this vertex. Indeed, if X is already a vertex of the odd cycle constructed above, then we are done. Otherwise the required cycle has three parts. The first part is a path from X to any vertex Y of the odd cycle, the second is the odd cycle, the third is the reverse path from Y to X .

Now we show that any two vertices can be connected by an odd path (not necessarily simple). Indeed, take two arbitrary vertices and connect them with an arbitrary path. This path is either odd or even. If it is odd, we are done. If it is even, we can combine it with an odd cycle passing through the first vertex of this path.

Now we explain how to obtain $(1, 1, 1, 0, 0, 0)$. This vector corresponds to the pair $(2, 3)$ of vertices of the graph. If they are already connected by an edge, we are done. Otherwise connect them with an odd path and take the alternating sum of its edges.

To finish the proof, explain how to obtain an arbitrary vector with the sum of coordinates ≥ 3 . We can obtain $(0, 1, 1, 1, 0, 0)$ in the same way as we obtain $(1, 1, 1, 0, 0, 0)$. Hence we can easily obtain $(1, 0, 0, -1, 0, 0)$ (and all the other vectors which can be obtained from this by a permutation of coordinates) as their difference. To get the required decomposition for an arbitrary vector, we will successively subtract vectors of form $e_i - e_j$ from our vector. At each step choose i and j such that the i th coordinate of our vector is maximal and the j th coordinate is minimal. If the difference between the maximal and the minimal coordinates is ≥ 2 , then in several steps it will diminish. If it equals 1, then the vector is equal to $(e_i + e_j + e_k) + a(1, 1, 1, 1, 1, 1)$, $a \in \mathbb{Z}$, but this vector equals $(e_i + e_j + e_k) \in \mathbb{Z}(v_1, \dots, v_5)$. If this difference equals 0, then the remaining vector equals 0. In both cases we are done. $\square$

Reformulate Lemma 2.5:

Lemma 2.6. *In the notation above, for any m linearly independent vectors $\{v_1, \dots, v_m\} \subseteq M(\lambda)$ the following equality holds:*

$$\mathbb{Z}(v_1, \dots, v_m) = \mathbb{Q}(v_1, \dots, v_m) \cap \mathbb{Z}_{\equiv 0(3)}(e_1, \dots, e_6).$$

Proof. If $m < 5$, we add several vectors from $M(\lambda)$ to this set to get a set of 5 linearly independent vectors $v_1, \dots, v_5$. Apply Lemma 2.5 to the set $v_1, \dots, v_5$. Since a vector has a unique representation on a basis,

$$\mathbb{Z}(v_1, \dots, v_m) = \langle v_1, \dots, v_m \rangle \cap \mathbb{Z}(v_1, \dots, v_5).$$

Finally, using Lemma 2.5, we get that $\mathbb{Z}(v_1, \dots, v_5)$ coincides with $\mathbb{Z}_{\equiv 0(3)}\{e_1, \dots, e_6\}$. $\square$

Remark. One may suppose that if we omit $(*)$ and take linearly independent vectors $v_1, \dots, v_6 \in \mathbb{Q}^6$, $v_i = e_p + e_q + e_r$, then $\mathbb{Z}(v_1, \dots, v_6) = \mathbb{Z}_{\equiv 0(3)}(e_1, \dots, e_6)$. However, this is not true – if we take the following vectors, then the volume of the unit cube will be equal to 6, and the index of the new lattice in $\mathbb{Z}^6$ will be 6, not 3.

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \\ v_6 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \end{pmatrix}, \quad \det \begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \\ v_6 \end{pmatrix} = 6.$$

Now we have to show that every subset $\{v_i\}$ in $M(\lambda)$ is saturated. To the contrary, let $\{v_1, \dots, v_m\} \subset M(\lambda)$ be an NSS. Then there exists a vector v and (by Lemma 2.3 and after renumbering) a linearly independent subset $\{v_1, \dots, v_s\} \subseteq \{v_1, \dots, v_m\}$ such that

$$v = q_1 v_1 + \dots + q_s v_s = z_1 v_1 + \dots + z_m v_m, \quad q_i \in \mathbb{Q}_+, z_i \in \mathbb{Z}.$$

This gives $v \in \mathbb{Z}_{\equiv 0(3)}(e_1, \dots, e_6)$, and $v \in \langle v_1, \dots, v_s \rangle$. Using Lemma 2.6, we obtain that $v \in \mathbb{Z}(v_1, \dots, v_s)$, $v = z'_1 v_1 + \dots + z'_s v_s$. Since $v_1, \dots, v_s$ are linearly independent, for any i we have $q_i = z'_i \in \mathbb{Z}_+$, and $v \in \mathbb{Z}_+(v_1, \dots, v_s)$. This shows that each subset of $M(\lambda)$ is saturated.

3. NEGATIVE RESULTS

Let λ be a highest weight not listed in the Main Theorem. One has to construct an NSS in $M(\lambda)$. There are two opportunities for λ : either the absolute values of all its usual coordinates are < 1 , or λ has a coordinate with the absolute value ≥ 1 . Speaking informally, the second case is practically always the consequence of the first one (Lemma 3.3), but the NSS in the first case is constructed recursively and its capacity increases when n increases. The construction of the second case gives an NSS of only 4 vectors for any n .

To prove that a set $\{v_1, \dots, v_m\}$ is not saturated, we will construct a so-called *discriminating function* $f(v)$ with the following properties: linearity, $= 0$ when $v = e_1 + \dots + e_n$ (to make it correctly defined), and non-negativity on the vectors of the set $\{v_1, \dots, v_m\}$. The discriminating function will be applied as follows. If we want to show that $\{v_0; v_1, \dots, v_m\}$ is an ENSS, it suffices to present the corresponding $\mathbb{Q}_+$ - and $\mathbb{Z}$ -combinations for v_0 and construct a discriminating function f , such that $f(v_0)$ cannot be composed as the sum of $f(v_i)$ with $\mathbb{Z}_+$ -coefficients.

Further, x_i denotes the function of taking the i th coordinate of a vector in some quasi-basis representation.

3.1. The fundamental weights. In this case λ equals

$$\pi_k = \pi_{k,n} = \left(\frac{n-k}{n}, \dots, \frac{n-k}{n}, -\frac{k}{n}, \dots, -\frac{k}{n} \right)$$

in the usual basis, $0 < k < n$, $n \geq 3$ (when $n = 2$, the corresponding representation is mentioned in the Main Theorem). In some proofs we will consider π_k for $SL(n)$'s of different dimensions simultaneously, so the second index in the notation $\pi_{k,n}$ carries this data. Here $M(\lambda) = \{\sigma\lambda \mid \sigma \in S_n\}$. The highest weight is equal to $e_1 + e_2 + \dots + e_k$ in quasi-basis, all the points of $M(\lambda)$ have a form $e_{i_1} + e_{i_2} + \dots + e_{i_k}$, $1 \leq i_1 < i_2 < \dots < i_k \leq n$.

Now we can reformulate the problem. Let $\{e_i\}$ be the quasi-basis, $k < n$, the weight $\lambda = \pi_k$ is not listed in the Main Theorem. One has to find a non-saturated subset in the set

$$\{e_{i_1} + e_{i_2} + \dots + e_{i_k} \mid 1 \leq i_1 < i_2 < \dots < i_k \leq n\}.$$


The construction will use induction on n . In the next section we will produce the NSSes which will be the base of the induction.

3.1.1. Important particular cases.

Example 1. $n = 7, k = 2$. The NSS will consist of those and only those vectors which are the sums of two quasi-basis vectors connected with an edge in the graph below. We have

$$v = e_1 + e_2 + e_3 = \frac{1}{2} \left((e_1 + e_2) + (e_2 + e_3) + (e_1 + e_3) \right),$$

$$v = -(e_4 + e_5 + e_6 + e_7) = 2(e_2 + e_3) - (e_2 + e_4) - (e_2 + e_5) - (e_3 + e_6) - (e_3 + e_7).$$



Let $f = 5(x_2 + x_3) - 2(x_1 + x_4 + x_5 + x_6 + x_7)$. Then

$$f(e_2 + e_3) = 10,$$

$$f(e_1 + e_2) = f(e_1 + e_3) = f(e_2 + e_4) = f(e_2 + e_5) = f(e_3 + e_6) = f(e_3 + e_7) = 3,$$

$$f(v) = f(e_1 + e_2 + e_3) = 5 \cdot 2 - 2 = 8.$$

It is clear that 8 cannot be represented as a sum where each summand equals either 3 or 10.

Example 2. $n = 8, k = 3$. Consider the following vectors (in quasi-basis):

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \\ v_6 \\ v_7 \\ v_8 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

Take $v = (1, 1, 1, 1, 1, 0, 0, 0) = \frac{1}{3}(v_1 + v_2 + v_3 + v_4 + v_5) = 2v_5 - v_6 - v_7 - v_8$. Let $f =$

$x_1 + 5(x_2 + x_3 + x_4) + 2x_5 - 6(x_6 + x_7 + x_8)$. Then

$$\begin{aligned} f(v_1) &= 12, & f(v_2) &= f(v_3) = 8, & f(v_4) &= 11, \\ f(v_5) &= 15, & f(v_6) &= f(v_7) = f(v_8) = 4, & f(v) &= 18. \end{aligned}$$

It is easy to see that 18 cannot be represented as a sum where each summand equals 4, 8, 11, 12, or 15.

Example 3. $n = 2k, k \geq 4$.

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ \vdots \\ v_{k-1} \\ v_k \\ v_{k+1} \\ v_{k+2} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 0 & 0 & 1 & 1 & \dots & 1 & 1 \\ 0 & 1 & 0 & \dots & 0 & 0 & 1 & 0 & 1 & \dots & 1 & 1 \\ 0 & 0 & 1 & \dots & 0 & 0 & 1 & 1 & 0 & \dots & 1 & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 & 1 & 1 & 1 & \dots & 0 & 1 \\ 0 & 0 & 0 & \dots & 0 & 1 & 1 & 1 & 1 & \dots & 1 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 & 1 & 1 & 1 & \dots & 1 & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 & 0 & 1 & 1 & \dots & 1 & 0 \end{pmatrix}$$

Show that it is an NSS. Let $v = (\underbrace{0, \dots, 0}_k, \underbrace{1, \dots, 1}_k)$, $v = v_1 + v_{k+1} - v_{k+2}$,

$$\begin{aligned} \frac{1}{k-2}(v_1 + \dots + v_k) &= \frac{1}{k-2}(\underbrace{1, \dots, 1}_k, \underbrace{k-1, \dots, k-1}_k) = \\ &= \frac{1}{k-2}(\underbrace{0, \dots, 0}_k, \underbrace{k-2, \dots, k-2}_k) = (\underbrace{0, \dots, 0}_k, \underbrace{1, \dots, 1}_k) = v. \end{aligned}$$

To explain why v is not a $\mathbb{Z}_+$ -combination of vectors v_i , consider two cases.

First case, when $k = 4$, let $f = -6x_3 - 7x_4 + 5(x_5 + x_6 + x_7) - 2x_8$. Then

$$f(v_1) = f(v_2) = 8, f(v_3) = 2, f(v_4) = 8, f(v_5) = 15, f(v_6) = 10, f(v) = 13.$$

But it is easy to see that 13 cannot be represented as a sum where each summand equals 2, 8, 10, or 15.

Second case, when $k \geq 5$, let $f = (k-2)(x_{k+1} + \dots + x_{2k}) - k(x_3 + \dots + x_k)$. Then

$$\begin{aligned} f(v_1) &= f(v_2) = (k-2)(k-1), \\ f(v_3) &= f(v_4) = \dots = f(v_k) = (k-1)(k-2) - k, \\ f(v_{k+1}) &= (k-2)(k-1), \\ f(v_{k+2}) &= (k-2)^2, \\ f(v) &= k(k-2). \end{aligned}$$

If $k \geq 6$, then two least possible summands give too much: $2((k-1)(k-2) - k) > k(k-2)$, if $k = 5$, then 15 should be represented as a sum where each summand equals 12, 7, or 9, but this is impossible.

Example 4. $n = 2k + 1, k \geq 3$.

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ \vdots \\ v_k \\ v_{k+1} \\ v_{k+2} \\ v_{k+3} \\ \vdots \\ v_{2k} \\ v_{2k+1} \end{pmatrix} = \begin{pmatrix} 0 & 1 & 1 & \dots & 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 1 & \dots & 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & \dots & 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & 1 & \dots & 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 1 & \dots & 1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 1 & 1 & 0 & 1 & 0 & \dots & 0 & 0 \\ 1 & 0 & \dots & 1 & 1 & 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & \dots & 0 & 1 & 0 & 0 & 0 & \dots & 1 & 0 \\ 1 & 1 & \dots & 1 & 0 & 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}$$

Let $v = (\underbrace{1, \dots, 1}_{k+1}, \underbrace{0, \dots, 0}_k)$. Then

$$v = \frac{1}{k}(v_1 + \dots + v_{k+1}) = \frac{1}{k}(\underbrace{k, \dots, k}_{k+1}, \underbrace{0, \dots, 0}_k) = (\underbrace{1, \dots, 1}_{k+1}, \underbrace{0, \dots, 0}_k),$$


$$\begin{aligned} (k-1)v_{k+1} - v_{k+2} - \dots - v_{2k+1} &= \\ &= (k-1)(\underbrace{1, \dots, 1}_k, \underbrace{0, \dots, 0}_{k+1}) - (\underbrace{k-1, \dots, k-1}_k, \underbrace{0, 1, \dots, 1}_k) = \\ &= (\underbrace{0, \dots, 0}_{k+1}, \underbrace{-1, \dots, -1}_k) = (\underbrace{1, \dots, 1}_{k+1}, \underbrace{0, \dots, 0}_k) = v. \end{aligned}$$

It suffices to show that v does not belong to $\mathbb{Z}_+(v_1, v_2, \dots, v_{2k+1})$. Let $f = (k+1)(x_1 + \dots + x_k) - k(x_{k+1} + \dots + x_{2k+1})$. Then

$$\begin{aligned} f(v_1) &= \dots = f(v_k) = k^2 - k - 1, \\ f(v_{k+1}) &= k(k+1), \\ f(v_{k+2}) &= \dots = f(v_{2k+1}) = k^2 - k - 1, \\ f(v) &= k^2. \end{aligned}$$

But if $k \geq 3$, then $k^2 < 2(k^2 - k - 1)$, so k^2 cannot be represented as a sum where each summand equals either $(k^2 - k - 1)$ or $k(k+1)$. This means that $v \notin \mathbb{Z}_+(v_1, v_2, \dots, v_{2k+1})$.

Example 5. $n = 8, k = 2$.



The NSS will contain those and only those vectors which are sums of two quasi-basis vectors connected with an edge in the graph above. Let $v = e_1 + e_2 + e_3 + e_5 + e_6 + e_7$. Then

$$v = \frac{1}{2}((e_1 + e_2) + (e_2 + e_3) + (e_1 + e_3) + (e_5 + e_6) + (e_6 + e_7) + (e_5 + e_7)),$$

$$v = (e_1 + e_2) + (e_3 + e_4) - (e_4 + e_5) + (e_5 + e_6) + (e_5 + e_7).$$

Check that $e_1 + e_2 + e_3 + e_5 + e_6 + e_7$ cannot be represented as a $\mathbb{Z}_+$ -combination of the vectors of our set. Let $f = x_1 + x_2 + x_3 + 2(x_5 + x_6 + x_7) + 9x_4 - 18x_8$. Then

$$\begin{aligned} f(e_1 + e_2) &= f(e_2 + e_3) = f(e_1 + e_3) = 2, & f(e_5 + e_6) &= f(e_6 + e_7) = f(e_5 + e_7) = 4, \\ f(e_3 + e_4) &= 10, & f(e_4 + e_5) &= 11, & f(v) &= 9. \end{aligned}$$

But 9 cannot be represented as the sum of integers 2, 4, 10, or 11.

Example 6. $n = 9, k = 3$.

Consider the following vectors:

$$\begin{aligned}
v_1 &= e_1 + e_2 + e_4, \\
v_2 &= e_1 + e_2 + e_5, \\
v_3 &= e_2 + e_3 + e_6, \\
v_4 &= e_2 + e_3 + e_7, \\
v_5 &= e_1 + e_3 + e_8, \\
v_6 &= e_1 + e_3 + e_9, \\
v_7 &= e_2 + e_4 + e_6.
\end{aligned}$$

Then $v = e_1 + e_2 + e_3 = \frac{1}{3}(v_1 + v_2 + v_3 + v_4 + v_5 + v_6) = v_1 + v_3 - v_7$.

Check that v is not a $\mathbb{Z}_+$ -combination of $v_1, v_2, v_3, v_4, v_5, v_6$, and v_7 . Let $f = 5(x_1 + x_2 + x_3 + x_4) - 4(x_5 + x_6 + x_7 + x_8 + x_9)$. Then

$$\begin{aligned}
f(v_1) &= 15, \\
f(v_2) &= f(v_3) = f(v_4) = f(v_5) = f(v_6) = f(v_7) = 6, \\
f(v) &= 15.
\end{aligned}$$

Note that $v \neq v_1$ and $f(v_1) = f(v)$, so we conclude that if $v \in \mathbb{Z}_+(v_1, \dots, v_7)$ then v_1 does not occur in this decomposition. But $15 \not\equiv 6$, this means that v cannot be obtained as a $\mathbb{Z}_+$ -combination of v_i 's.

Example 7. $n = 10, k = 4$. Consider the following vectors:

$$\begin{aligned}
v_1 &= e_1 + e_2 + e_3 + e_5, \\
v_2 &= e_1 + e_2 + e_4 + e_6, \\
v_3 &= e_3 + e_4 + e_5 + e_6, \\
v_4 &= e_5 + e_6 + e_7 + e_8, \\
v_5 &= e_5 + e_7 + e_8 + e_9, \\
v_6 &= e_6 + e_7 + e_8 + e_{10},
\end{aligned}$$

$$v = e_1 + e_2 + e_3 + e_4 + e_5 + e_6 = \frac{1}{2}(v_1 + v_2 + v_3) = v_4 - v_5 - v_6.$$

Show that $v \notin \mathbb{Z}_+(v_1, \dots, v_6)$. Let $f = x_1 + x_3 + x_4 + 6x_7 + 6x_8 - 7x_9 - 8x_{10}$. Then

$$\begin{aligned}
f(v_1) &= f(v_2) = f(v_3) = 2, \\
f(v_4) &= 12, f(v_5) = 5, f(v_6) = 4, \\
f(v) &= 3.
\end{aligned}$$

But it is clear that 3 cannot be represented as a sum where each summand equals 2, 4, 5, or 12.

3.1.2. Case when $n \not\equiv k, n \not\equiv (n-k)$.

It follows from these two conditions that $n \geq 5$. The exceptional case $\frac{k}{n} \in \{\frac{2}{5}, \frac{3}{5}\}$ will be considered at the end of the section. Further we (temporarily) suppose that $\frac{k}{n} \notin \{\frac{2}{5}, \frac{3}{5}\}$, which gives $n \geq 7$.

Lemma 3.1. *Suppose that there exists an NSS for a pair (n, k) , where (n, k) satisfy the conditions above. Then for each $r \in \mathbb{N}$, there exists an NSS for the pair (nr, kr) .*

Proof. Consider an arbitrary vector from $M(\pi_{k,n})$. Write down its quasi-coordinates r times in succession. The result is a vector from $M(\pi_{kr,nr})$: it has kr 1's and $(n-k)r$ 0's. If one takes an NSS for (n, k) and performs this procedure on each vector, the result will be an NSS for (nr, kr) . $\square$

Thus, if we construct an NSS for all pairs (n, k) where $\gcd(n, k) = 1$, then the NSS for all other pairs will be also constructed according to the Lemma.

Lemma 3.2 (the Step procedure). *Having constructed an NSS for a pair (n, k) , one can construct an NSS for the pair $(n+k, k)$ according to the existing NSS.*

Proof. The keypoint is that if one takes a weight from $M(\pi_{k,n})$, writes it down in the form where all its quasi-coordinates are equal to 0 or to 1, and adds k coordinates equal to 0, then this weight can be considered as a weight from $M(\pi_{k,n+k})$. If we start with an ENSS $(v; v_1, v_2, \dots, v_m)$ for (n, k) , we should perform this procedure on all its vectors and then add one more vector v_{m+1} which has 0s at the first n digits and 1s at the k adjoint digits. Now we show that the obtained set $(v'; v'_1, \dots, v'_m, v_{m+1})$ is indeed an NSS in $M(\pi_{k,n+k})$.

Suppose that a vector v lies in the ENSS for (n, k) , $v = q_1 v_1 + q_2 v_2 + \dots + q_s v_s$, $q_i \in \mathbb{Q}_+$. If we fix some representations for v and for all v_i in quasi-basis, then this equality can be re-written in the *formal* basis in the following form:

$$v = q_1 v_1 + q_2 v_2 + \dots + q_s v_s - \alpha(f_1 + f_2 + \dots + f_n),$$

where f_i 's are the counter images of e_i 's under the projection $\mathbb{Q}^n \rightarrow \mathfrak{X}(T) \otimes_{\mathbb{Z}} \mathbb{Q}$, $(q_1, q_2, \dots, q_n) \mapsto q_1 e_1 + q_2 e_2 + \dots + q_n e_n$. Obviously, f_i 's are linearly independent. For each v_i , fix a representation in which it has k coordinates equal to 1 and $n-k$ coordinates equal to 0. The vector v is nonzero, consequently, it has a representation where all its coordinates are nonnegative, but some of them are zeroes. Fix this representation. Then $\alpha \geq 0$ (otherwise all coordinates of v are strictly positive), and we get that in $\mathbb{Q}^{n+k}$ the following equality holds:

$$v' = q_1 v'_1 + q_2 v'_2 + \dots + q_s v'_s + \alpha(f_{n+1} + \dots + f_{n+k}) - \alpha(f_1 + f_2 + \dots + f_{n+k}).$$

This shows that v' lies in the $\mathbb{Q}_+$ -lattice generated by $v'_1, \dots, v'_m, v_{m+1}$ (here all vectors taken in quasi-basis $\{e_1, \dots, e_{n+k}\}$).

Similarly one can show that v' still lies in the $\mathbb{Z}(v'_1, \dots, v'_m, v_{m+1})$.

To prove that the constructed set is indeed an ENSS, it remains to show that v' does not lie in $\mathbb{Z}_+(v'_1, \dots, v'_m, v_{m+1})$. Suppose the contrary. Let $v' \in \mathbb{Z}_+(v'_1, \dots, v'_m, v_{m+1})$. Omit last k coordinates. We get that $v \in \mathbb{Z}_+(v_1, \dots, v_m)$, so $\{v; v_1, v_2, \dots, v_m\}$ is not an ENSS for (n, k) . $\square$

Now we can explain how, using these Lemmas, the NSS's can be constructed for all pairs (n, k) , for which the following three conditions are held:

- (1) $1 < k < n - 1$,
- (2) $\gcd(n, k) = 1$,
- (3) $n \geq 7$.

Use descent on n . Suppose the NSSes are constructed for all pairs (m, l) , satisfying the conditions above, with $m < n$. Take a pair (n, k) . Suppose $k < \frac{n}{2}$ (otherwise change it by $n - k$ and seek for an NSS for the pair $(n, n - k)$, the case $n = 2k$ is impossible because $\gcd(n, k) = 1$). If all the conditions are held for the pair $(n - k, k)$, then we have an NSS for it, and using the Step procedure, this NSS can be re-made into the NSS for (n, k) . Let us find all the cases when at least one of the conditions fails for $(n - k, k)$.

Condition (1) fails iff $n = 2k + 1$. But we have $n \geq 7$, then $k \geq 3$. In this case we already have an NSS (example 4).

Condition (2) never fails.

Condition (3) fails iff $n - k \leq 5$. Find these cases. Recall that $k \leq (n - 1)/2$. Substitute it: $n \leq (n - 1)/2 + 5$. This gives $n \leq 9$. List all these pairs (n, k) (with $k < \frac{n}{2}$).

$n = 7$. Pairs $(7, 2)$ and $(7, 3)$.

$n = 8$. Pair $(8, 3)$.

$n = 9$. Pairs $(9, 2)$ and $(9, 4)$.

But we already have NSSes for all these pairs. Indeed, cases $(7, 2)$ and $(8, 3)$ coincide with Examples 1 and 2, respectively. Cases $(7, 3)$ and $(9, 4)$ are the particular cases of $n = 2k + 1$ (Example 4). Case $(9, 2)$ can be obtained from $(7, 2)$ (Example 1) using the Step procedure.

Finally, take all the cases where the NSS is already constructed as the base of the descent. In all the other cases the descent is feasible, consequently, we have constructed an NSS for all pairs (n, k) for which the conditions (1) – (3) hold.

Now we consider the case $\frac{k}{n} \in \{\frac{2}{5}, \frac{3}{5}\}$. Let $k = 2k_1$, $n = 5k_1$, $k_1 \geq 2$. When $k_1 \geq 4$, we can construct an NSS using the Step procedure and substitution $k \rightarrow n - k$: starting with an NSS for $(2k_1, k_1)$, we successively construct NSSes for $(3k_1, k_1)$, $(3k_1, 2k_1)$, and $(5k_1, 2k_1)$. When $k_1 = 2$, the Example 7 can be applied.

When $k_1 = 3$, the pair $(n, k) = (15, 6)$, and the required NSS can be obtained from Example 6 using the Step procedure.

3.1.3. Case when $n \vdots k$ or $n \vdots (n - k)$.

Assume that $k \leq n/2$. Then $n \vdots k$, $n = kd$. In the case when $k = 1$ all the subsets in the sets of weights are saturated (see 2.1), further $k \geq 2$.

When $k \geq 4$, Example 3 shows that the NSS exists for the pair $(2k, k)$. Using the Step procedure, we can easily rebuild this NSS into the NSS for a pair (kd, d) , where $d \geq 2$. It remains to consider cases $k = 2$ and 3.

$k = 2$. It follows from the Main Theorem that $d \geq 4$. But we already have an NSS for the pair $(8, 2)$ (Example 5), using the Step procedure, we can construct NSSes for all $d > 4$.

$k = 3$. We already have an NSS for $(9, 3)$, using the Step procedure, we can construct an NSS for all $n \vdots 3$, $n > 9$.

We are done.

3.2. Non-fundamental weights. In the previous section the structure of $M(\lambda)$ was much easier than in the general case. Indeed, by definition $M(\lambda) = (\lambda + \Phi) \cap P$, where $P = \text{conv}\{\sigma\lambda \mid \sigma \in S_n\}$. If λ is a fundamental weight, then we just take $\{\sigma\lambda \mid \sigma \in S_n\}$, not dealing with Φ , because it does not add new points. However, if λ is not fundamental, then $M(\lambda)$ contains internal points of P , and this is a great advantage for constructing NSSes.

Lemma 3.3 (Inclusion Lemma). *Let λ and λ' be two dominant weights, such that $\lambda' \in M(\lambda)$, and there exists an NSS in $M(\lambda')$. Then there exists an NSS in $M(\lambda)$.*

Proof. Notice that $\forall \sigma \in W \quad \sigma\lambda' \in M(\lambda)$ and $P' = \text{conv}\{\sigma\lambda' \mid \sigma \in W\} \subset P$. Then $M(\lambda') = (\lambda' + \Phi) \cap P' \subset M(\lambda)$. This means that the NSS for λ' is also an NSS for λ . $\square$

There are two cases: the first one – all the usual coordinates of λ are integer, and the second one – all of them are non-integer. Consider these cases independently. The coordinates of vectors in the usual basis will be denoted by y_i . The discriminating functions in the usual basis must satisfy only the conditions of linearity and $f(v_i) \geq 0$.

3.2.1. All the coordinates of λ are integer, $\lambda \neq (1, 0, \dots, 0, -1)$.

Definition. By a *Shift* we will denote the following procedure: take a point $\lambda = (y_1, \dots, y_n)$, fix two indices $i < j$ such that $|y_i - y_j| \geq 2$ and replace λ with the point λ' , where $\lambda' = (\dots, y_i - 1, \dots, y_j + 1, \dots)$ if $y_i > y_j$ and $(\dots, y_i + 1, \dots, y_j - 1, \dots)$ otherwise.

The point λ' lies in $M(\lambda)$. Indeed, $M(\lambda)$ contains the point $(\dots, y_j, \dots, y_i, \dots)$, its convex hull with λ (with the proper coefficient) contains λ' . Notice that after each Shift $y_1^2 + \dots + y_n^2$ diminishes by a positive integer. Indeed, let $x = \max\{y_i, y_j\}$, $y = \min\{y_i, y_j\}$, then $x - y \geq 2$,

$$(x - 1)^2 + (y + 1)^2 = x^2 - 2x + 1 + y^2 + 2y + 1 = x^2 + y^2 - 2(x - y - 1) \leq x^2 + y^2 - 2.$$

This means that if we apply consequent Shifts to λ , then this process cannot be infinite.

Lemma 3.4. *If $n \geq 3$ and λ satisfies the conditions of subsection 3.2.1, then $M(\lambda)$ contains one of the points $(2, 0, \dots, 0, -1, -1)$, $(1, 1, 0, \dots, 0, -2)$, or $(1, 1, 0, \dots, 0, -1, -1)$, and it always contains the point $(1, 0, \dots, 0, -1)$.*

Proof. Let $\lambda = (a_1, \dots, a_n)$ (in the usual basis). If $\forall i \ a_i \in \{-1, 0, 1\}$, then, due to the fact that $\lambda \neq (1, 0, \dots, 0, -1)$, λ has at least 4 nonzero coordinates. Taking into account that $\sum_1^n a_i = 0$, at least two of them are equal to 1 and two are equal to -1 . In this case, $M(\lambda)$ contains the point $(1, 1, 0, \dots, 0, -1, -1)$: split all its other coordinates into pairs 1, -1 and make them zero (using the Shift), then permute the remaining 4 coordinates. Applying one more Shift, we yield $(1, 0, \dots, 0, -1)$.

Otherwise, if $\exists i, |a_i| > 1$ (one of the coordinates is *big*), then $\max_{i,j} (a_i - a_j) \geq 3$. Keeping at least one coordinate big, perform the shift for the pairs of indices where $|a_i - a_j| \geq 2$. This process is finite. Consider a situation where we can perform no more Shift. If we still have a nonzero coordinate with the same sign as the big coordinate has, we can shift it with the coordinate of the opposite sign (their difference will obviously be ≥ 2). Otherwise we are in the case where we have a big coordinate of one sign (without loss of generality positive) and some coordinates of the opposite sign. If the big coordinate is ≥ 3 , then apply a Shift

to this coordinate and to some negative coordinate. But we have supposed that Shifts are impossible. Then the big coordinate is equal to 2, nonzero ones among the other coordinates are either -2 or -1 and -1 . But if $M(\lambda)$ contains a point $(2, 0, 0, \dots, 0, -2)$, then it also contains $(2, 0, \dots, 0, -1, -1) = \frac{1}{2}((2, 0, \dots, 0, -2) + (2, 0, \dots, 0, -2, 0))$. We can easily get $(1, 0, \dots, 0, -1)$, performing one more Shift. $\square$

Construct NSSes for the first three points.

Example 8. $\lambda = (2, 0, \dots, 0, -1, -1)$, $n \geq 3$.

Consider vectors

$$\begin{aligned} v_1 &= (1, -1, 0, 0, \dots, 0), \\ v_2 &= (-1, -1, 2, 0, \dots, 0), \\ v_3 &= (2, -1, -1, 0, \dots, 0), \\ v &= (0, -1, 1, 0, \dots, 0) = \frac{1}{2}(v_1 + v_2) = v_2 + v_3 - v_1. \end{aligned}$$

Suppose $f = -y_2$, then $f(v_1) = f(v_2) = f(v_3) = f(v) = 1$, but $\forall i \ v \neq v_i$. We get a contradiction.

The NSS for the point $\lambda = (1, 1, 0, \dots, 0, -2)$, $n \geq 3$ can be constructed similarly (one should multiply all the coordinates by -1).

Example 9. $\lambda = (1, 1, 0, \dots, 0, -1, -1) \in M(\lambda)$, $n \geq 4$.

Consider vectors

$$\begin{aligned} v_1 &= (1, 1, -1, -1, 0, \dots, 0), \\ v_2 &= (1, -1, 1, -1, 0, \dots, 0), \\ v_3 &= (0, 1, 0, -1, 0, \dots, 0), \\ v_4 &= (0, 0, 1, -1, 0, \dots, 0), \\ v &= (1, 0, 0, -1, 0, \dots, 0) = \frac{1}{2}(v_1 + v_2) = v_1 + v_4 - v_3. \end{aligned}$$

Suppose $f = -y_4$, then $f(v_1) = f(v_2) = f(v_3) = f(v_4) = f(v) = 1$, but $v \neq v_i$ for any i . We get a contradiction.

Now take an arbitrary dominant weight λ , $n \geq 3$, and the corresponding set $M(\lambda)$. It follows from Lemma 3.4 and the Inclusion Lemma that the NSS for λ exists.

It remains to consider the case $n = 2$, $\lambda = (a, -a)$. If $|a| \geq 3$, then $\text{conv}\{(\sigma(a_1, a_2)), \sigma \in S_2\}$ contains the points $(2, -2)$, and $(3, -3)$. But this subset is not saturated:

$$(1, -1) = \frac{1}{2}(2, -2) = (3, -3) - (2, -2),$$

and the vector $(1, -1)$ is not a linear combination of vectors $(2, -2)$ and $(3, -3)$ with integer positive coefficients. If, otherwise, $a \in \{0, \pm 1, \pm 2\}$, then each subset in $M(\lambda)$ is saturated (sections 2.2 and 2.4).

3.2.2. All the coordinates of λ are non-integer.

Lemma 3.5. *Given a point $\lambda = (a_1, a_2, \dots, a_n)$ (in the usual basis), $n \geq 4$. If the set $\{a_1, a_2, \dots, a_n\}$ contains simultaneously $\alpha + 1, \alpha, \alpha - 1$ for some $\alpha \in \mathbb{R}$, then the set $M(\lambda)$ contains an NSS.*

Proof. It is easy to see that $M(\lambda)$ contains a point $v_1 = (\alpha + 1, \alpha, \alpha - 1, a_4, \dots, a_n)$, $a_4 \neq 0$ (because $a_4 \notin \mathbb{Z}$). Acting by S_n , we can get the following points from it:

$$\begin{aligned} v_2 &= (\alpha - 1, \alpha, \alpha + 1, a_4, \dots, a_n), \\ v_3 &= (\alpha + 1, \alpha - 1, \alpha, a_4, \dots, a_n), \\ v_4 &= (\alpha, \alpha - 1, \alpha + 1, a_4, \dots, a_n). \end{aligned}$$

Show that this set is not saturated. Indeed,

$$\begin{aligned} \frac{1}{2}(v_1 + v_2) &= (\alpha, \alpha, \alpha, a_4, \dots, a_n), \\ v_2 + v_3 - v_4 &= (\alpha, \alpha, \alpha, a_4, \dots, a_n), \end{aligned}$$

suppose $f = \frac{y_4}{a_4}$, then

$$f(v_1) = f(v_2) = f(v_3) = f(v_4) = f(v) = 1.$$

But $\forall i \quad v \neq v_i$, this means that v is not a $\mathbb{Z}_+$ -combination of v_i . □

Lemma 3.6 (Good Triple Lemma). *Let $\lambda = (a_1, \dots, a_n)$, $n \geq 4$, and all a_i are non-integer. If the collection $a_1, \dots, a_n$ contains at least three different values, then $M(\lambda)$ contains a point of form $(\alpha + 1, \alpha, \alpha - 1, a_4, \dots, a_n)$.*

Proof. Perform several Shifts preserving the condition that the set $\{a_1, \dots, a_n\}$ contains at least 3 elements. Suppose further Shifts are impossible (we mentioned above that, starting from any position, only a finite number of Shifts is possible). Consider $a_{\max} = \max\{a_1, \dots, a_n\}$, $a_{\min} = \min\{a_1, \dots, a_n\}$, $a_{\text{mid}} \in \{a_1, \dots, a_n\}$, $a_{\text{mid}} \neq a_{\max}$, $a_{\text{mid}} \neq a_{\min}$. If $a_{\max} - a_{\text{mid}} \geq 3$, then we can apply the Shift to $a_{\max}$ and a_{mid} , thus we obtain three different values of coordinates $a_{\min}$, $a_{\text{mid}} + 1$, $a_{\max} - 1$. Similarly, if $a_{\text{mid}} - a_{\min} \geq 3$, then at least one more Shift is possible. So we yield $a_{\max} - a_{\text{mid}}, a_{\text{mid}} - a_{\min} \in \{1, 2\}$. If $a_{\max} - a_{\text{mid}} = a_{\text{mid}} - a_{\min} = 1$, we have already found a point of necessary type in $M(\lambda)$. Up to symmetry, one of the two cases is possible: either $a_{\min} = a_{\text{mid}} - 2$, $a_{\max} = a_{\text{mid}} + 2$, or $a_{\min} = a_{\text{mid}} - 1$, $a_{\max} = a_{\text{mid}} + 2$. Consider these two cases.

In the first case, apply the Shift to $a_{\max}$ and $a_{\min}$. This operation gives us the required triple $(a_{\max} - 1, a_{\text{mid}}, a_{\min} + 1)$.

In the second case, $a_{\min} = a_{\text{mid}} - 1$, $a_{\max} = a_{\text{mid}} + 2$, and we know that λ has at least 4 coordinates. If there are 4 different values among them, the fourth will inevitably form a triple of form $(\alpha + 1, \alpha, \alpha - 1)$ with two of $a_{\max}$, a_{mid} , $a_{\min}$. Otherwise $a_i \in \{a_{\max}, a_{\text{mid}}, a_{\min}\}$ for any i . But $n \geq 4$, this means that at least one of the values $(a_1, a_2, \dots, a_n)$ is mentioned twice. Suppose $n = 4$ (we need only 4 a_i 's, forget that there are other coordinates). The multiplicities of $(a_{\max}, a_{\text{mid}}, a_{\min})$ may be as follows: $(\widehat{1}, \widehat{1}, 2)$, $(\widehat{1}, 2, 1)$, $(2, 1, 1)$. Apply the shift to the coordinates marked with the hat. We get one of the following collections: $(a_{\text{mid}} + 1, a_{\text{mid}}, a_{\text{mid}}, a_{\text{mid}} - 1)$, $(a_{\text{mid}} + 1, a_{\text{mid}} + 1, a_{\text{mid}}, a_{\text{mid}} - 1)$, $(a_{\text{mid}} + 2, a_{\text{mid}} + 1, a_{\text{mid}}, a_{\text{mid}})$. Each of them contains a triple of form $(\alpha + 1, \alpha, \alpha - 1)$. But this means that here we also find a triple of form $(\alpha + 1, \alpha, \alpha - 1)$. □

Lemma 3.7 (Absence of a good triple). *Let $\lambda = (a_1, \dots, a_n)$ be a dominant weight, $n \geq 4$, $\exists i$ with $|a_i| > 1$, and all $a_i \notin \mathbb{Z}$. If $M(\lambda)$ does not contain a point of the form $(\alpha + 1, \alpha, \alpha - 1, a_4, \dots, a_n)$, then*

$$\lambda = \left(\frac{2n-2}{n}, -\frac{2}{n}, -\frac{2}{n}, \dots, -\frac{2}{n} \right) \text{ or } \lambda = \left(\frac{2}{n}, \frac{2}{n}, \dots, \frac{2}{n}, -\frac{2n-2}{n} \right).$$

Proof. If the collection $a_1, \dots, a_n$ contains at least 3 different elements, then we can use the Good Triple Lemma and show that $M(\lambda)$ contains a point of the desired form. This means that $\forall i, a_i \in \{a_{\max}, a_{\min}\}$. Without loss of generality, we may suppose that $a_{\max} > 1$, and $a_{\min} < 0$ (otherwise multiply all a_i by -1).

If $a_{\min} < -1$, then apply the Shift to $a_{\min}$ and $a_{\max}$. Thus we get $a_{\min} + 1$ and $a_{\max} - 1$ among the values of the coordinates, and still at least one of $a_{\min}$ and $a_{\max}$ is presented (since $n \geq 4 > 3$). Using the Good Triple Lemma, we get a contradiction.

We see that $-1 < a_{\min} < 0$. If the collection $(a_1, \dots, a_n)$ contains $a_{\max}$ at least for 2 times, then apply the shift to $a_{\min}$ and $a_{\max}$. Now we have $a_{\max}, a_{\min} + 1 > 0$ and at least one time $a_{\min}$ among the values of coordinates: all the coordinates cannot be positive. This gives us a contradiction with the Good Triple Lemma.

Then $a_{\max}$ enters only once in $(a_1, \dots, a_n)$. If $a_{\max} > 2$, apply the Shift to $a_{\min}$ and $a_{\max}$. We get that $a_{\max} - 1 > 1$, $a_{\min} + 1 < 1$ and $a_{\min}$ are among the values of coordinates, which gives us a contradiction with the Good Triple Lemma.

We yield that the collection has a form $(a_{\max}, a_{\min}, a_{\min}, \dots, a_{\min})$, $1 < a_{\max} < 2$, $-1 < a_{\min} < 0$. Let $a_{\min} = -\frac{k}{n}$. We have $(n-1)a_{\min} + a_{\max} = 0$ from the initial conditions. This yields $a_{\max} = \frac{k(n-1)}{n}$. But $a_{\max} < 2$. Consequently,

$$\frac{k(n-1)}{n} < 2 \Rightarrow (n-1) < 2n \Rightarrow k < \frac{2n}{n-1} < 3,$$

because $n \geq 4$. Taking into account that $a_{\max} > 1$, we get $k = 2$, $a_{\max} = \frac{2n-2}{n}$, $a_{\min} = -\frac{2}{n}$. But in the beginning of the case we could change the signs at all the coordinates. Thus, we have two cases: $\lambda = (\frac{2n-2}{n}, -\frac{2}{n}, -\frac{2}{n}, \dots, -\frac{2}{n})$ and $\lambda = (-\frac{2n-2}{n}, \frac{2}{n}, \frac{2}{n}, \dots, \frac{2}{n})$. $\square$

Applying the Lemmas, we see that in the case, when $a_i \notin \mathbb{Z}$, $\exists i, |a_i| > 1$, $n \geq 4$, we have not constructed an NSS only in these two cases. In all other cases the NSS exists due to Lemma 3.5. Let us construct an NSS in these two cases. We may assume that $\lambda = (\frac{2n-2}{n}, -\frac{2}{n}, -\frac{2}{n}, \dots, -\frac{2}{n}) = 2e_1$. Let

$$v_1 = 2e_1,$$

$$v_2 = 2e_2,$$

$$w = 2e_3,$$

$$v_3 = e_1 + e_3 \in M(\lambda),$$

$$v_4 = e_2 + e_3 \in M(\lambda).$$

Then v_1, v_2, v_3, v_4 form an NSS. Indeed, we have

$$v = e_1 + e_2 = \frac{1}{2}(v_1 + v_2) = v_1 + v_4 - v_3,$$

$$f = x_1 + x_2 + x_3 - 3x_n.$$

Then $f(v_1) = f(v_2) = f(v_3) = f(v_4) = f(v) = 2$, and $v \neq v_i$ for any i . But 2 cannot be represented as a sum of more than one 2s. We get a contradiction.

Now it remains to consider the cases $n = 3$ and $n = 2$.

In the case $n = 3$ we suppose that the fractional parts of all coordinates are equal to $\frac{2}{3}$ (otherwise change λ for $-\lambda$, as we've done earlier). If $\lambda = (\frac{2}{3}, \frac{2}{3}, -\frac{4}{3})$, then each subset in $M(\lambda)$ is saturated (see 2.6). Below we construct an NSS for $\lambda = (\frac{5}{3}, -\frac{1}{3}, -\frac{4}{3}) = 3e_1 + e_2$, then, using the Inclusion Lemma, show the existence of NSS for all other points λ . Let

$$\begin{aligned} v_1 &= e_1 = \frac{2}{3}(3e_1 + e_2) + \frac{1}{3}(e_2 + 3e_3), \\ v_2 &= 2e_1 + 2e_2 = \frac{1}{2}(3e_1 + e_2) + \frac{1}{2}(e_1 + 3e_2), \\ v_3 &= 3e_1 + e_2, \\ v &= 2e_1 + e_2 = v_1 + \frac{1}{2}v_2 = v_3 - v_1. \end{aligned}$$

If $f = x_1 - x_3$, then $f(v_1) = 1$, $f(v_2) = 2$, $f(v_3) = 3$, and $f(v) = 2$. But v is equal neither to v_2 , nor to $2v_1$. We get a contradiction.

Lemma 3.8. *Suppose that $\lambda = (a_1, a_2, a_3)$ (in the usual basis) is a dominant weight such that the fractional parts of all a_i are equal to $\frac{2}{3}$. Suppose also that there exists an index i with $|a_i| > 1$, and $\lambda \neq (\frac{2}{3}, \frac{2}{3}, -\frac{4}{3})$. Then $M(\lambda)$ contains a point $(\frac{5}{3}, -\frac{4}{3}, -\frac{1}{3})$.*

Proof. It follows from the conditions of the lemma that $\exists i, a_i \geq \frac{5}{3}$. Indeed, otherwise we have at least 2 positive coordinates, each of them $\leq \frac{2}{3}$, but due to the condition of the Lemma there exists an a_i such that $|a_i| > 1$. Suppose it is a_1 . We have $a_1 = -a_2 - a_3 \geq -\frac{4}{3}$. This means that $\lambda = (-\frac{4}{3}, \frac{2}{3}, \frac{2}{3})$. We get a contradiction.

If only one coordinate of λ is positive, and it is equal to $\frac{5}{3}$, then $\lambda = (\frac{5}{3}, -\frac{4}{3}, -\frac{1}{3})$, and the Lemma is proved. Otherwise either λ has two positive coordinates, or one of them is $\geq \frac{8}{3}$. In both cases we can apply the Shift to a positive and a negative coordinate, such that after it λ still has a coordinate $\geq \frac{5}{3}$, and so on. $\square$

Consider the case $n = 2$. Suppose $\lambda = (\frac{a}{2}, -\frac{a}{2})$, $a \in \mathbb{N}$, a is odd. Then for $a = 3$ each subset in $M(\lambda)$ is saturated (see 2.3). When $a \geq 5$, we construct an NSS. Let

$$v_1 = \left(\frac{3}{2}, -\frac{3}{2}\right), \quad v_2 = \left(\frac{5}{2}, -\frac{5}{2}\right).$$

Then $(\frac{1}{2}, -\frac{1}{2}) = \frac{1}{3}v_1 = 2v_2 - v_1 \notin \mathbb{Z}_+(v_1, v_2)$.

So we have constructed non-saturated subsets in the sets of weights for all the representations not listed in the Main Theorem.

REFERENCES

- [BZ] G. Bobiński, G. Zwara, Normality of orbit closures for directing modules over tame algebras, J. Algebra **298** (2006), 120–133
- [Chi] C. Chindris, Orbit Semigroups and the Representation Type of Quivers, <http://arxiv.org/abs/0708.3413v1>
- [CK] J. B. Carrell, A. Kurth, Normality of Torus Orbit Closures in G/P , J. Algebra **233** (2000), 122–134

- [Da] R. Dabrowski, On Normality of the Closure of a Generic Torus Orbit in G/P , Pacific Journal of Mathematics, **192**, No. 2 (1996), 321–330
- [Ful] W. Fulton, Introduction to Toric Varieties, Princeton University Press, 1993
- [Ha] F. Harary, Graph Theory, Addison-Wesley Publishing Company, 1969
- [Hu] J. E. Humphreys, Introduction to Lie Algebras and Representation Theory, GTM, **9**, Springer-Verlag, 1978
- [KKMSD] G. Kempf, F. Knudsen, D. Mumford, B. Saint-Donat, Toroidal Embeddings I, LNM, **339**, Springer-Verlag, Berlin-Heidelberg-New York, 1973
- [Kl] A. A. Klyachko, Toric Varieties and Flag Varieties, Tr. Mat. Inst. Steklova, **208** (1995), 139–162. English translation: A. A. Klyachko, Toric Varieties and Flag Varieties, Proc. Steklov Inst. Math., **208** (1995), 124–145
- [Mo] J. Morand, Closures of torus orbits in adjoint representations of semisimple groups, C.R.Acad.Sci Paris Sér.I Math. **328**, No.3 (1999), 197–202
- [OH] H. Ohsugi and T. Hibi, Normal polytopes arising for finite graphs, J. Algebra **207** (1998), 409–426
- [Stu1] B. Sturmfels, Equations Defining Toric Varieties, Proc. Sympos. Pure Math., **62**, Part 2, AMS, Providence, RI, 1997, 437–449
- [Stu2] B. Sturmfels, Gröbner Bases and Convex Polytopes, University Lecture Series, **8**, AMS, Providence, RI, 1996
- [SVV] A. Simis, W. Vasconcelos, and R. Villarreal, The integral closure of subrings associated to graphs, J. Algebra **199** (1998), 281–299
- [VO] Vinberg, E.B., Onishchik, A.L., Seminar on Lie groups and algebraic groups, Moskva, Nauka, 1988 (Russian). English translation: A.L. Onishchik, E.B. Vinberg, Lie groups and algebraic groups, Berlin Heidelberg New York, Springer 1990
- [Wh] N.White, The basis monomial ring of a matroid, Advances in Math. **24** (1977), 292–297

DEPARTMENT OF HIGHER ALGEBRA, FACULTY OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, 119992 MOSCOW, RUSSIA

E-mail address: karina@mccme.ru